



THE SPECTRAL PROBLEM OF THE LAPLACE OPERATOR ON A CIRCLE

Bobojonov Jurabek

Mathematics Master's Student, Asian International University

Abstract. *This paper investigates the spectral problem of the Laplace operator defined on a circle. The problem is formulated on a one-dimensional compact smooth manifold under periodic boundary conditions. It is shown that the Laplace operator is a self-adjoint and positive operator with a purely discrete spectrum. The eigenvalues are explicitly determined and form an unbounded increasing sequence, while the corresponding eigenfunctions are represented by trigonometric functions. These eigenfunctions constitute a complete orthonormal basis in the Hilbert space $L^2(S^1)$ establishing a direct connection between the spectral decomposition of the Laplace operator and classical Fourier analysis. In addition, the asymptotic behavior of the eigenvalues is analyzed and shown to be consistent with general spectral laws for compact manifolds. The results obtained provide a clear and rigorous framework for understanding fundamental spectral properties and have important applications in mathematical physics, particularly in quantum mechanics and heat conduction models.*

Keywords. *Laplace operator; spectral problem; circle; eigenvalues; eigenfunctions; Fourier analysis; periodic boundary conditions; spectral theory.*

Introduction. Spectral theory of differential operators plays a fundamental role in modern mathematical physics, functional analysis, and geometry. Among these operators, the Laplace operator occupies a central position due to its wide applicability in describing physical phenomena such as heat conduction, wave propagation, quantum mechanics, and vibration analysis. The investigation of

spectral properties of the Laplace operator on manifolds with simple geometry provides essential insight into more complex spectral problems.

The circle represents one of the simplest compact one-dimensional manifolds without boundary, making it an ideal model for studying the spectral problem of the Laplace operator. Despite its apparent simplicity, the Laplace operator on a circle exhibits rich mathematical structure, closely connected with Fourier analysis, eigenvalue distributions, and periodic boundary conditions. The explicit form of eigenfunctions and eigenvalues allows for a complete and rigorous spectral characterization, which serves as a benchmark for understanding spectral behavior on higher-dimensional or more intricate manifolds.

The spectral problem of the Laplace operator on a circle is formulated as an eigenvalue problem for a second-order differential operator with periodic boundary conditions. Its solutions are closely related to trigonometric functions, and the resulting spectrum is discrete and unbounded. These properties make the circle an important testing ground for theoretical concepts such as spectral asymptotics, eigenvalue multiplicity, and completeness of eigenfunction systems.

Moreover, studying the Laplace operator on a circle has significant implications beyond pure mathematics. In physics, it models quantum particles constrained to move on a ring, circular membranes, and periodic media. In engineering and signal processing, the spectral decomposition on a circle underlies harmonic analysis techniques used in data representation and filtering.

The aim of this paper is to analyze the spectral problem of the Laplace operator on a circle, focusing on the formulation of the eigenvalue problem, the derivation of eigenvalues and eigenfunctions, and the investigation of their fundamental properties. By providing a clear and systematic exposition of this classical problem, the paper contributes to a deeper understanding of spectral theory on compact manifolds and establishes a foundation for further generalizations and applications.

Literature Review. The spectral theory of the Laplace operator has been extensively studied in classical and modern mathematical literature. Foundational results in this field were established within the framework of functional analysis and partial differential equations, where the Laplace operator serves as a prototypical example of a self-adjoint elliptic operator on compact manifolds. Early systematic treatments of eigenvalue problems for second-order differential operators can be found in the seminal works of Richard Courant and David Hilbert, who laid the theoretical foundations of spectral analysis and variational methods. Their contributions clarified the relationship between boundary conditions, eigenvalue distributions, and orthogonality of eigenfunctions [1,2].

A comprehensive functional-analytic approach to the spectral problem of the Laplace operator is presented in the works of Michael Reed and Barry Simon, where self-adjoint operators, spectral decomposition, and unbounded operators in Hilbert spaces are studied in depth. These works emphasize the role of compactness and periodic boundary conditions, which are directly relevant to the Laplace operator on a circle. The explicit solvability of the problem on one-dimensional compact manifolds is often used as an illustrative example for general spectral theorems.

From a geometric perspective, the spectral properties of the Laplace operator on compact manifolds, including the circle, have been analyzed within spectral geometry. Classical results demonstrate how the geometry and topology of a manifold influence the structure of the spectrum. In this context, the circle serves as a fundamental example where the spectrum can be computed explicitly, allowing for a clear illustration of abstract geometric principles. Studies in this area highlight the discreteness of the spectrum and the symmetry-induced multiplicity of eigenvalues.

The connection between the Laplace operator on a circle and Fourier analysis is well documented in harmonic analysis literature. Periodic functions on a circle naturally lead to Fourier series expansions, where eigenfunctions of the Laplace operator form a complete orthonormal basis in appropriate function spaces. This relationship has been explored in numerous textbooks and research articles,

emphasizing the importance of trigonometric eigenfunctions in solving differential equations with periodic conditions.

In applied mathematics and physics, the spectral problem of the Laplace operator on a circle has been investigated in relation to quantum mechanics, wave propagation, and heat transfer in periodic or circular domains. The model of a quantum particle on a ring, for instance, relies directly on the spectral properties of the Laplacian under periodic boundary conditions. Such applications demonstrate that the theoretical results obtained for the circle are not merely abstract but have direct physical interpretations.

Overall, the existing literature provides a solid theoretical and applied foundation for the study of the spectral problem of the Laplace operator on a circle. However, revisiting this classical problem remains valuable, as it offers a clear and rigorous setting for illustrating fundamental concepts of spectral theory and serves as a reference point for more complex generalizations to higher-dimensional or nontrivial geometries.

Main Part. 1. Mathematical Formulation of the Problem.

Let S^1 denote a circle of length L , parametrized by the interval $[0, L]$ with periodic identification of endpoints. The Laplace operator is defined as:

$$\Delta u(x) = d^2u(x)/dx^2$$

The spectral problem is formulated as:

$$-d^2u(x)/dx^2 = \lambda u(x),$$

with periodic boundary conditions:

$$u(0) = u(L), u'(0) = u'(L).$$

2. Self-Adjointness and Spectral Properties

The operator $A = -d^2/dx^2$ acting on the Hilbert space $L^2(S^1)$ is self-adjoint, positive, and has a discrete spectrum.

Its eigenvalues are real, non-negative, and have finite multiplicity. The eigenfunctions form a complete orthonormal basis in $L^2(S^1)$.

3. Eigenvalues and Eigenfunctions

For $\lambda = 0$, the solution is a constant function, hence $\lambda_0 = 0$.

For $\lambda > 0$, solutions take the form:

$$u_n(x) = \cos(2\pi nx/L), \quad v_n(x) = \sin(2\pi nx/L),$$

with eigenvalues:

$$\lambda_n = (2\pi n/L)^2, \quad n \in \mathbb{Z}.$$

Each nonzero eigenvalue has multiplicity two.

4. Orthogonality and Completeness

The eigenfunctions constitute an orthonormal basis in $L^2(S^1)$. Any function $f \in L^2(S^1)$ can be expanded into a Fourier series using these eigenfunctions.

5. Spectral Asymptotics

The eigenvalues satisfy the asymptotic relation:

$$\lambda_n \sim (2\pi n/L)^2 \text{ as } n \rightarrow \infty,$$

which is consistent with Weyl-type asymptotic laws for compact manifolds.

6. Physical Interpretation

The model describes a quantum particle on a ring and heat propagation in circular media. Eigenvalues correspond to energy levels, while eigenfunctions describe stationary states.

Conclusion. In this paper, the spectral problem of the Laplace operator on a circle has been investigated in a rigorous and systematic manner. By formulating the

problem on a one-dimensional compact manifold with periodic boundary conditions, the study demonstrates how fundamental concepts of spectral theory can be explicitly realized in a simple geometric setting.

The analysis shows that the Laplace operator on a circle is a self-adjoint, positive operator with a purely discrete spectrum. The eigenvalues are explicitly determined and form an unbounded sequence, while the corresponding eigenfunctions are given by trigonometric functions. The multiplicity of nonzero eigenvalues reflects the rotational symmetry of the circle, highlighting the close connection between geometry and spectral properties.

It has been established that the system of eigenfunctions constitutes a complete orthonormal basis in the Hilbert space $L^2(S^1)$. As a result, any square-integrable function on the circle admits a Fourier series representation, confirming the equivalence between the spectral decomposition of the Laplace operator and classical Fourier analysis. This property plays a crucial role in both theoretical investigations and practical applications.

Furthermore, the asymptotic behavior of the eigenvalues is consistent with general spectral laws for compact manifolds, illustrating how the circle serves as a fundamental example for understanding more complex spectral phenomena. The explicit solvability of the problem provides valuable insight into the structure of eigenvalue distributions and serves as a reference model for higher-dimensional or more intricate geometries.

In conclusion, the spectral problem of the Laplace operator on a circle not only offers a complete and transparent mathematical framework but also forms a conceptual bridge between abstract spectral theory and applied fields such as quantum mechanics, heat conduction, and signal processing. The results obtained in this study may serve as a foundation for further generalizations, including perturbations of the operator, non-uniform metrics, or extensions to higher-dimensional compact manifolds.

References

1. Courant, R., & Hilbert, D. (1989). *Methods of Mathematical Physics. Vol. I.* New York: Wiley-Interscience.
2. Courant, R., & Hilbert, D. (1989). *Methods of Mathematical Physics. Vol. II.* New York: Wiley-Interscience.
3. Reed, M., & Simon, B. (1980). *Methods of Modern Mathematical Physics. Vol. I: Functional Analysis.* New York: Academic Press.
4. Davies, E. B. (1995). *Spectral Theory and Differential Operators.* Cambridge: Cambridge University Press.
5. Berger, M. (2003). *A Panoramic View of Riemannian Geometry.* Berlin: Springer.
6. Zygmund, A. (2002). *Trigonometric Series.* Cambridge: Cambridge University Press.
7. Evans, L. C. (2010). *Partial Differential Equations.* Providence, RI: American Mathematical Society.
8. Griffiths, D. J. (2018). *Introduction to Quantum Mechanics.* Cambridge: Cambridge University Press.